



Contribution ID: 49

Type: Paper

TO STUDY THE LEVEL OF IRISIN IN PATIENTS WITH PCOS

Saturday, 18 September 2021 17:25 (10 minutes)

Background –

PCOS is a complex condition with reproductive, endocrine and metabolic characteristics. For its diagnosis, Rotterdam Criteria is used : 2 of 3 criteria should be present-

- Oligomenorrhea/ Amenorrhea
- Biochemical or clinical signs of hyperandrogenism
- Polycystic ovaries with >20 follicles in ovary or ovarian volume >10cc

****Irisin**** is a newly discovered peptide hormone , a cleavage product of fibronectin type III domain 5

Aims & objectives-

- To determine Serum Irisin levels in cases with PCOS and healthy controls
- To compare Serum Irisin levels in cases with PCOS and healthy controls

Setting: Dr BRAMH, Raipur , OPD patients

Method- Total 90 patients, out of this 30 Obese PCOS patients, 30 non-obese PCOS patients and 30 healthy controls. Participants 2ml fasting blood sample was taken for estimation of Irisin ,by commercially available ELISA kit . Anthropometric measurements taken and hormonal profile also evaluated.

Results- In our study Polycystic ovary syndrome women showed different body compositions compared with controls(non- PCOS). Serum irisin level of PCOS did not show significant difference compared with controls.Irisin maybe raised in obese PCOS patients as compared to non obese PCOS.

Primary authors: VERMA, Kritika (Post graduate 3rd year); Dr GUPTA, Ruchi (Pt. JNM Medical College, Raipur); Dr NAGARIA, Tripti (Pt. JNM Medical College, Raipur)

Presenter: VERMA, Kritika (Post graduate 3rd year)

Session Classification: Paper Presentation Slot 4