



Contribution ID: 15

Type: **Paper**

## A rare case of Vaginal Leiomyoma

Leiomyoma are benign smooth muscle tumors commonly occurring in the uterus and to some extent in the cervix followed by the round ligament, utero-sacral ligament, ovary, and inguinal canal. Vaginal leiomyoma are very rare with only about 300 reported cases. A 49 yrs old para 3 lady presented in the gynaecological outpatient department with complains of lower abdominal pain, polymenorrhagia, dysmenorrhoea, dyspareunia, urgency and frequency of micturition since 6 months. Medical and surgical history were not significant. On per speculum examination the vagina was occupied with a large mass arising from the anterior vaginal wall and cervix could not be visualised. On bimanual examination, a firm globular smooth surfaced mass 6x5 cm was felt arising from the middle third of anterior vaginal wall up to 3 cm from the introitus with restricted mobility and non-tender. MRI pelvis reported as diffuse adenomyosis and a well defined rounded mass lesion arising from anterior vaginal wall displacing the base of uterus anteriorly suggestive of vaginal leiomyoma. Transvaginal removal of mass by enucleation along with total abdominal hysterectomy with bilateral salpingo-oophorectomy was done and postoperative diagnosis was anterior vaginal wall leiomyoma which was supported by histopathology report.

**Primary authors:** Dr AGRAWAL, Ankita (MS Obstetrics & Gynaecology); Dr CHATURVEDI, Srishti (DGO, Secondary DNB OBGY Resident)

**Presenters:** Dr AGRAWAL, Ankita (MS Obstetrics & Gynaecology); Dr CHATURVEDI, Srishti (DGO, Secondary DNB OBGY Resident)